Contents Volume 15, 1990

Late Neogene calcareous nannofossil biostratigraphy of the Gulf of Aden region	
A. Rahman and P.H. Roth (Salt Lake City, UT, U.S.A.)	1
Planktonic-benthonic foraminiferal ratios: Modern patterns and Tertiary applicability	
T.G. Gibson (Reston, VA, U.S.A.)	29
Distribution of diatoms in sediments of the northern Indian Ocean: relationship to physical oceanography L.H. Burckle (Palisades, NY, U.S.A.)	53
Relation of counting methods to taphonomic gradients and biofacies zonation of foraminiferal sediment assemblages R.E. Martin (Newark, DE, U.S.A.) and W.D. Liddell (Logan, UT, U.S.A.)	
Multiple Miocene marine productivity events in West Antarctica as recorded in upper Miocene sediments beneath the Ross Ice Shelf (Site J-9)	
D.M. Harwood, R.P. Scherer and PN. Webb (Columbus, OH, U.S.A.)	01
Syndepositional dissolution of calcareous foraminifera in modern shallow-water sediments	31
J.W. Murray (Exeter, England)	117
Seasonal variation in boreal Early Cretaceous calcareous nannofossils	111
E. Thomsen (Aarhus, Denmark)	193
Palaeoceanography of the last deglaciation in the Alboran Sea (western Mediterranean). Stable isotopes and	120
planktonic foraminiferal records	150
C. Pujol (Talence, France) and C. Vergnaud Grazzini (Paris, France)	153
Modern and late Quaternary pollen spectra of fiord sediments, eastern Baffin Island, Arctic Canada	101
S.K. Short (Boulder, CO, U.S.A.), J.T. Andrews (Boulder, CO, U.S.A.) and W.N. Mode (Oshkosh, WI, U.S.A.) Ostracoda in the Earth Sciences by P. de Decker, JP. Colin and JP. Peypouquet — H.J. Oertli	181
Introduction	203
J.R. Blueford (Fremont, CA, U.S.A.)	205
Distribution, abundance and diversity of Antarctic acantharian cysts	200
M. Spindler and K. Beyer (Bremerhaven, F.R.G.)	200
Comparing radiolarian and diatom diversity and abundance from the northeast Pacific	203
J.R. Blueford (Fremont, CA, U.S.A.), J.J. Gonzales (San José, CA, U.S.A.) and K. Van Scoy	
(Miami, FL, U.S.A.)	219
Experimental evidence for changes during fossilization of radiolarian tests and implications for a model of	210
biomineralization	
M.S. Afanasieva (Moscow, U.S.S.R.)	233
Radiolarian petrograhic preservation types in Jurassic to Lower Tertiary cherts of Costa Rica	200
HJ. Gursky (Marburg/Lahn, F.R.G.)	249
Evolutionary trends of Antarctissa in the Quaternary using morphometric analysis	
A. Granlund (Stockholm, Sweden)	265
Origin of the subgenera Cyclampterium, Paralampterium and Sciadiopeplus from Lophocytris (Lophocytris)	
(Radiolaria, Theoperidae)	
A. Sanfilippo (La Jolla, CA, U.S.A.)	287
New biostratigraphic elements for dating upper Norian strata from the Sandilands Formation, Queen Charlotte	
Island, British Columbia, Canada	
E.S. Carter (Veronia, OR, U.S.A.)	313
Middle to Late Jurassic radiolarians from Sicily (Italy)	
N. Kito, P. De Wever, T. Danelian and F. Cordey (Paris, France)	329
Evolutionary trends and biostratigraphic potential of selected radiolarian taxa from the Early Carboniferous of	
Germany	
A. Braun (Marburg/Lahn, F.R.G.)	351
Significance of Devonian-Carboniferous radiolarians from accretionary terranes of the New England Orogen, eastern Australia	
J. Aitchison (Armidale, Australia)	365

Paleoenvironmental and biostratigraphic significance of siliceous microfossils of the Permo-Triassic	Redding Section,
Eastern Klamath Mountains, California	
P. Noble (Austin, TX, U.S.A.) and P. Renne (Princeton, NJ, U.S.A.)	379
Ordovician Radiolaria from Nevada and Newfoundland — A comparison at the family level	
G.W. Renz (La Jolla, CA, U.S.A.)	393

()

